

## SEI Drop Regimen by Surgeon 6/5/24

Doctor	Drop or injection	Surgery Type	Pre-Op Drops	Pre-Op Instructions	Post-Op Drops	Post-Op Instructions	Notes / Substitutes
Brown	TriMoxi Injection	Cataract	N/A		N/A	N/A	*Majority of patients get Injection is for all locations *If patient using drops; same as Venerable. (i.e. Trabec or some allergies)
Brown	Drops	LASIK			1. Prednisolone Acetate / Gatifloxacin combo gtt 2. Artificial Tears prn	1. Use 1 gtt in surgical eye(s) QID x 1 week 2. Use artificial tears PRN	
Brown	Drops	PRK			1. Prednisolone Acetate / Gatifloxacin / Bromfenac combo gtt 2. Artificial Tears prn	1. Use 1 gtt in surgical eye(s) QID until bottle gone 2. Use artificial tears PRN	
Chai	Drops	Cataract	N/A	N/A	1. Prolensa 2. Pred Forte 3. Moxifloxacin	1. 1 drop once a day for 30 days 2. 1 drop four times a day for 14 days, then 1 drop 2 times a day for 14 days (28 days total) 3. 1 drop four times a day for 7 days	
Chai	Drops	Cataract			Pred-Moxi-Brom (Compound drop )	1 drop three times in the operative eye for 30 days; start day of surgery	
Cherry	Drops	Cataract	Polytrim/Ocuflox		1. Polytrim/Ocuflox 2. Predforte	1. 1 drop 4x per day for 7 days 2. 1 drop 4x per day for 7 days, then 1 drop 3x per day for 7 days, then 1 drop 2x per day for 7 days, then 1 drop 1x per day for 7 days	
Cherry	TriMoxi Injection	Cataract	N/A	N/A	N/A	N/A	Injection is for all locations including Hickman Hospital and Adrian
Cherry	Drops	Glaucoma	Prolensa		1. Polytrim/Ocuflox 2. Predforte	1. 1 drop 4x per day for 7 days 2. 1 drop 4x per day for 7 days, then TBD depending on post-op appointments.	

El Yassir	Drop	Cataract	N/A	N/A	1. Prolensa 2. Pred Forte 3. Ofloxacin	1. 1 drop once a day for 30 days 2. 1 drop four times a day for 14 days, then 1 drop 2 times a day for 14 days (28 days total) 3. 1 drop four times a day for 7 days	
El Yassir	Drop	Cataract			Pred-Moxi-Brom (Compound drop )	1 drop three times in the operative eye for 30 days; start day of surgery	
Ernest	Drop	Cataract	Ocuflox/Prolensa	1 drop the night before and morning of surgery	1. Prolensa 2. Pred Forte 3. Ofloxacin	1. 1 drop once a day for 30 days 2. 1 drop four times a day for 30 days 3. 1 drop four times a day for 7 days	
Ernest	Drop	Cataract (SAME DAY SX)	Ocuflox/Prolensa		1. Prolensa 2. Pred Forte 3. Ofloxacin	1. 1 drop once a day for 30 days 2. 1 drop four times a day for 30 days 3. 1 drop four times a day for 7 days	
Ernest	Drop	Cataract			1. Pred-Moxi-Brom (Compound drop ) 2. then Pred -Brom (Compound drop )	(Pred-Moxi-Brom) 1 drop three times in the operative eye for 7 days (Pred-Brom) 3 times a day in the operative eye for 21 days	Substitute: Pred QID x 30 days, Ocuflox QID x 7 days and Illevro QD x 30 days Allergies: NKDA or list of allergies (If allergies apply type this at the end: "(per doctor, ok to fill)" – Dr. Ernest wants the patient to try the drop w/ punctal occlusion if history of allergies.)
Ernest	Drop	Cataract (SAME DAY SX)			1. Pred-Moxi-Brom (Compound drop ) 2. then Pred -Brom (Compound drop )	(Pred-Moxi-Brom) 1 drop three times in the operative eye for 7 days (Pred-Brom) 3 times a day in the operative eye for 21 days	Substitute: Pred QID x 30 days, Ocuflox QID x 7 days and Illevro QD x 30 days Allergies: NKDA or list of allergies (If allergies apply type this at the end: "(per doctor, ok to fill)" – Dr. Ernest wants the patient to try the drop w/ punctal occlusion if history of allergies.)
Ernest	Drops	PTK and PRK	None	None	1.Ocuflox 2. Pred Forte 3.Illevro/Prolensa 4.Preservative Free Systane	1. One drop in the operative eye four times a day for 7 days 2. One drop four times a day for 14 Days, then decrease to 1 drop 3 times a day for 7 days, then 1 drop 2 times a day for 7 days, then 1 drop a day for another 7 days, then stop 3. One drop daily for 3 days then stop 4. One drop every 2 hours while awake	

Gago	Drops	Cataract	Polytrim/Ocuflox	1 drop night before surgery and 1 drop morning of surgery	1. Polytrim/Ocuflox 2. Pred Forte 1%	1. Polytrim or Ocuflox (tan top, generic is ofloxacin) – 1 drop 4 times a day in the operative eye for 7 days. 3. Pred Forte – 1 drop 4 times a day in the operative eye for 30 days.	
Gago	TriMoxi Injection + Drop	Cataract			Pred- Brom ( <i>Compound drop</i> )	1. 1 drop in operative eye 3 times a day for 30 days	
Gago	Drops	LASIK	None	None	1. PF Artificial tears 2. Ocufloxacin 3. Prednisone Acetate	1. 1 drop every 2 hours while awake 2. 1 drop QID x 7 days 3. q 1 hr day of surgery and then QID x 7 days	
Gago	Drops	LASIK			Pred- Moxi ( <i>Compound drop</i> )	1. Use 1 drop every 2 hours while awake in the surgical eye(s) day of surgery. Then 1 drop 4 times a day for 7 days in the surgical eye(s)	
Gago	Drops	PRK/PTK	None	None	1. PF Artificial tears 2. Ocufloxacin 3. Prednisone Acetate	1. 1 drop every 2 hours while awake 2. 1 drop QID x 7 days 3. QID x 7 days	
Gago	Drops	EVO Implantable Contact Lens	None	None	1. Ocuflox 2. Pred-Brom (combo drop)		
Gordon	Drop	Retina	N/A	N/A	1. Prednisolone 2. Ocuflox	1. Prednisolone 1%- 1 drop in surgical eye QID times a day for 30 days 2. Ocuflox 0.3%- 1 drop in surgical eye QID times a day for 7 days	
Lavery	Drops	LASIK			1.Ocuflox 2. Pred Forte 3.Preservative Free Systane	1. One drop in the operative eye four times a day for 7 days 2. One drop four times a day for 7 days 3.One drop in the operative eye every 2 hours while awake	
Lavery	Drops	LASIK			Pred- Moxi ( <i>Compound drop</i> )	1. Use 1 drop every 2 hours while awake in the surgical eye(s) day of surgery. Then 1 drop 4 times a day for 7 days in the surgical eye(s)	
Lavery	Drops	PRK & PTK			Pred- Moxi ( <i>Compound drop</i> )	1. Use 1 drop every 2 hours while awake in the surgical eye(s) day of surgery. Then 1 drop 4 times a day until bottle is gone	
Lavery	Drops	PRK & PTK			1.Ocuflox	1. One drop in the operative eye four times a day for 7 days	
Lemon	Drop	Cataract	Besivance, Prolensa, Inveltys	Start Besivance 1 drop three times a day, Inveltys 1 drop two times a day, Prolensa 1 drop once a day. All start two days before surgery	1. Besivance 2. Inveltys 3. Prolensa	1. Besivance three times a day x 1 week 2. Inveltys twice a day for 28 days after 1 week appt 3. Prolensa once day for 28 days after 1 week appt	

Surendar Purohit	Drop	Retina	N/A	N/A	1. Prednisolone 2. Ocuflox	1. Prednisolone 1%- 1 drop in surgical eye QID times a day for 30 days 2. Ocuflox 0.3%- 1 drop in sx eye QID times a day for 7 days	
Ravin	Drop	Cataract	Ocuflox Ilevro / Prolensa	1 drop night before surgery and 1 drop morning of surgery	1. Ocuflox 2. Ilevro / Prolensa 3. Durezol	1. Ocuflox (tan top, generic is ofloxacin) – 1 drop 4 times a day in the operative eye. Use for 7 days after surgery. 2. Ilevro/Prolensa (no generic) – 1 drop once daily in the operative eye. Continue for 14 days after surgery. 3. Durzol(no generic) – 1 drop 2 times a day in the operative eye for 14 days.	Can replace Durezol with Preforte if too expensive or samples are not available at the office.
Rhem	Drop	Cataract	Ocuflox/Prolensa	1 drop the night before and morning of surgery	1.Ocuflox 2. Prolensa 3. Pred Forte	1. 1 drop 4x/day x 7 days 2. 1 drop once a day x 30 days 3. 1 drop 4 x/day x 14 days, then 1 drop 2x/day x 14 days (28 days total)	
Rhem	TriMoxi Injection + Drop	Cataract			Pred-Brom ( <i>Compound drop</i> )	1 drop two times in the operative eye for 30 days; start day of surgery	*Majority of patients DO get Trimoxi including glaucoma/cataract combo
Rhem	Drop	Cataract			Pred-Moxi-Brom ( <i>Compound drop</i> )	1 drop three times in the operative eye for 30 days; start day of surgery	
Sensoli	Drops	Cataract	Polytrim/Ocuflox	1 drop night before surgery and 1 drop morning of surgery	1. Polytrim/Ocuflox 2. Pred Forte 1%	1. Polytrim or Ocuflox (tan top, generic is ofloxacin) – 1 drop 4 times a day in the operative eye for 7 days. 3. Pred Forte – 1 drop 4 times a day in the operative eye for 30 days	NOT using Imprimis drops at this time.
Sensoli	Drops	LASIK			1. PF Artificial tears 2. Ocufloxacin 3. Prednisone Acetate	1. 1 drop every 2 hours while awake 2. 1 drop QID x 7 days 3. q 1 hr day of surgery and then QID x 7 days	NOT using Imprimis drops at this time.
Sensoli	Drops	LASIK			Pred- Moxi ( <i>Compound drop</i> )	1. Use 1 drop every 2 hours while awake in the surgical eye(s) day of surgery. Then 1 drop 4 times a day for 7 days in the surgical eye(s)	NOT using Imprimis drops at this time.
Sensoli	Drops	PRK/PTK			1. PF artificial tears 2. Ocufloxacin 3. Prednisone Acetate	1. 1 drop every 2 hours while awake 2. 1 drop QID x 7 days 3. QID x 7 days	NOT using Imprimis drops at this time.
Simpson	TriMoxi Injection + Drop	Cataract			Pred-Brom ( <i>Compound drop</i> )	1 drop two times in the operative eye for 30 days; start day of surgery	*Majority of patients DO get Trimoxi including glaucoma/cataract combo
Simpson	Drop	Cataract	Ocuflox/Prolensa	1 drop the night before and morning of surgery	1.Ocuflox 2. Prolensa 3. Pred Forte	1. 1 drop 4x/day x 7 days 2. 1 drop once a day x 30 days 3. 1 drop 4 x/day x 14 days, then 1 drop 2x/day x 14 days (28 days total)	

Simpson	Drop	Cataract			Pred-Moxi-Brom (Compound drop )	1 drop three times in the operative eye for 30 days; start day of surgery	
Simpson	Drops	LASIK			1. Prednisolone Acetate (Pred Forte 1%) 2. Ocufloxacin 3. Lubrication Drops	1. Use 1 drop every hour while awake in the surgical eye(s) day of surgery. Then 1 drop 4 times a day for 7 days in the surgical eye(s) 2. Use 1 drop four times a day for 7 days 2. Use 1 drop every 3. hours while awake x 1 week, then 4 times a day as needed	
Simpson	Drops	LASIK			Pred- Moxi (Compound drop )	1. Use 1 drop every 2 hours while awake in the surgical eye(s) day of surgery. Then 1 drop 4 times a day for 7 days in the surgical eye(s)	
Simpson	Drops & Pill Form	PRK			1. Prednisolone Acetate (Pred Forte 1%) 2. Ilevro 3. Ocufloxacin 4. Neurotin / Gabapentin 5. Lubrication Drops	1. Day of surgery, use on drop every hour while awake in the surgical eye(s), then 1 drop 4x a day for 14 days, 1 drop 3x a day for 7 days, 1 drop 2x a day for 7 days, 1 drop once per day for 7 days, then stop 2. 1 drop once a day x 3 days 3. Use 1 drop four times a day for 7 days 4. 1 tab 3 times a day by mouth for 3 days 5. 1 drop 4 times a day as needed	
Simpson	Drops	PRK			Pred-Moxi (Compound drop )	1. Use 1 drop every 2 hours while awake in the surgical eye(s) day of surgery. Then 1 drop 4 times a day for 14 days until bottle is gone	
Tolchin	Drop	Cataract	Polytrim/Ocuflox Prolensa	1 drop night before surgery and 1 drop morning of surgery	3. Lubrication Drops	1. Polytrim or Ocuflox (tan top, generic is ofloxacin) – 1 drop 4 times a day in the operative eye. Use for 7 days after surgery. 2. Prolensa– 1 drop once daily in the operative eye. Continue for 30 days after surgery. 3. Durezol- 1 drop 2 times daily in operative eye for 14 days after surgery.	Dr. Tolchin will sub out prolensa with either bromsite or illvero Can add will sub PRED FORTE 1% if Durezol is too expensive. Pred Forte instructions would be QID in operative eye for 14 days after surgery.
Tolchin	TriMoxi Injection	Cataract	NA	NA	NA	NA	*Majority of patients get Injection is for all locations including Hickman Hospital
Venerable	Drops	Cataract	Polytrim/Ocuflox Prolensa	1 drop night before surgery and 1 drop morning of surgery	1. Polytrim/Ocuflox 2. Durezol 3. Prolensa	1. Polytrim or Ocuflox (tan top, generic is ofloxacin) – 1 drop 4 times a day in the operative eye. Use for 7 days after surgery. 2. Prolensa– 1 drop once daily in the operative eye. Continue for 30 days after surgery. 3. Durezol- 1 drop 2 times daily in operative eye for 14 days after surgery.	Dr. Venerable will sub out prolensa with either bromsite or illvero Will substitute PRED FORTE 1% if Durezol is too expensive. Pred Forte instructions would be QID in operative eye for 14 days after surgery.

Venerable	TriMoxi Injection	Cataract	NA	NA	NA	NA	*Majority of patients get Injection is for all locations including Hickman Hospital
-----------	-------------------	----------	----	----	----	----	---